

TESTING KITS FOR DRUGS

Procedure Card for IDT2200

MIX HEROIN IDENTIFICATION

Substance Test



Read **Before** Use


As a precaution, gloves and eye protection should be worn when handling drug substances and using test units. Do not prove probe once a capsule is broken. In the unlikely event of a spill, rinse affected area thoroughly with water. Keep out of reach of children. All results are presumptive. If necessary, further testing by a competent drug testing lab can provide additional identification.



If the suspected substance is a brown powder, proceed to Steps 2-5 on Side 1.


If the suspected substance is a white powder or a tablet, proceed to Steps 6-11 on Side 2.

Brown Heroin Test





Gently shake test unit for 2-3 seconds.



Brown Heroin



If the substance is a brown powder, the appearance of a purple-red color strongly indicates the presence of Brown Heroin.



If no purple-red color develops, Brown Heroin is not present. Regardless of whether or not any color developed, do not continue to White Heroin testing. **STOP** and proceed to Disposal.



The test kit is safe to dispose of in the trash.



Manufactured by IDenta Ltd www.identa-corp.com Patented Technology • Made in Israel




Procedure Card for IDT2200

MIX HEROIN IDENTIFICATION

Substance Test

SIDE 2

White Heroin Test



With thumb and forefinger, squeeze chamber 2 until White Heroin Identification capsule breaks. DO NOT (2) 3 3 **BREAK CHAMBER 1**

Gently shake test unit for 2-3 seconds.



If the substance is a powder, the appearance of a purple color indicates the presence of an opiate such as Heroin.

If the substance is a tablet, the appearance of a purple color indicates the presence of an opiate such as Morphine or Codeine.



If no purple color develops, an opiate, such as Heroin or Morphine, is not present. Regardless of whether or not any color developed, do not continue to Brown Heroin testing. **STOP** and proceed to Dilution & Disposal.



Proceed to Dilution (steps 10-11), regardless of whether or not a color appeared.





Gently shake test unit for 2-3 seconds.



The test window will warm up. This is normal.

Disposal

After Dilution, the test kit is safe to dispose of in the trash.



Manufactured by IDenta Ltd www.identa-corp.com Patented Technology • Made in Israel